



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.


Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

# GEOMETRICAL DEMONSTRATION OF A THEOREM.

BY ISAAC H. TURRELL, CUMMINSVILLE, OHIO.

Let  $T$  be the common tangent, and  $d_1, d_3$  the diameters of the two circles that can be described touching three circles that touch each other, then  $T^2 = 4d_1d_3$ .



This theorem is given by Matthew Collins, on page 278 of *Mathematical Monthly*, Vol. 1, who states that he proposed it in an old number of the "Educational Times," but no geometrical demonstration had yet appeared. He then gives a solution of a particular case, only, namely, that in which one of the three circles that touch each other becomes infinite.

The following method pre-supposes some knowledge of the properties of Centers of Similitude, of radical axes and of circles in contact, and it is based on a beautiful theorem, remarkable for its generality, known to the ancients under the name of "The Arbelos," or "The Shoemaker's Knife," which is enunciated by Mr. Collins, in the paper referred to, by means of a figure, as follows:

"If the semicircles on the diameters  $AB, AC$ , touching each other at  $A$ , be both touched by the circles whose centers are  $O$  and  $O'$ , which touch each other at  $L$ ; demit  $OF, O'H$  perpendiculars on  $AB$ , then if  $OF = n$  times the diameter of  $O$ ,  $O'H = (n + 1)$  times the diameter of  $O'$ ,  $O'$  being nearer to  $A$  than  $O$  is."

The elegant geometrical proof there given, holds true, if the two original semicircles touched each other externally at  $A$ .

Let the circle whose center is  $O''$ , touch the circle  $O'$  and the two original semicircles,  $O'''$  touch  $O''$  and the semicircles, and so on in the same order,  $N$  being the center of the  $n^{\text{th}}$  circle from  $O$ ; it is required to find the relation connecting the quantities  $T, R$  and  $r$ ,  $T$  being the

common tangent, and  $R, r$ , the radii of the circles whose centers are  $O$  and  $N$ .

This may be easily accomplished by means of the Arbelos, thus:

Putting  $OF = m$  times  $2R$ , then  $NW = (m + n).2r$ .

$$\frac{NW}{2r} - \frac{OF}{2R} = n. \quad NW.R - OF.r = 2nRr \dots\dots\dots(1).$$

Putting  $ON = d$ ,  $d^2 = T^2 + (R - r)^2 \dots\dots\dots(2).$

Now produce  $ON$  until it intersects the common tangent, or radical axis of the original semicircles in  $P$ ; hence by a well-known theorem, namely, "If each of two circles touch (in the same way) another pair of circles, the center of similitude of either pair lies upon the radical axis of the other pair,"  $P$  is the external center of similitude of the circles  $O$  and  $N$ , and

$$\frac{PO}{PN} = \frac{R}{r} \dots\dots\dots(3).$$

Also  $S, S'$ , being two anti-homologous points, (see Chauvenet's, p. 360, or any treatise on the Modern Geometry) on the circles  $O$  and  $N$ , we have the equation

$$\sqrt{(PN + NS')(PO - OS)} = PA \dots\dots\dots(4).$$

From (3) we get  $\frac{dr}{R - r} = PN, \frac{dR}{R - r} = PO$ .

Hence (4) becomes

$$\sqrt{\left(\frac{dr}{R - r} + r\right)\left(\frac{dR}{R - r} - R\right)} = \sqrt{\frac{d^2 R r}{(R - r)^2} - R r} = PA \dots\dots\dots(5).$$

By similar triangles

$$\frac{PA - OF}{PA - NW} = \frac{PO}{PN} = \frac{R}{r}.$$

$$PA(R - r) = NW.R - OF.r = 2nRr \text{ from (1).}$$

$PA = \frac{2nRr}{R - r}$ . By substituting in eq. (5), this value of  $PA$ , and the value of  $d^2$  as given in (2), we get

$$\left\{ \frac{[T^2 + (R - r)^2]Rr}{(R - r)^2} - Rr \right\}^{\frac{1}{2}} = \frac{2nRr}{R - r},$$

which reduces to  $T = 2n\sqrt{Rr}$ , the relation required; or putting  $R = \frac{1}{2}d_1$ ,  $r = \frac{1}{2}d_n$ , we get  $T = n\sqrt{d_1d_n}$ .

If  $n = 2$ , we have the original theorem proposed by Mr. Collins.

The same method will apply, if the two original semicircles touched each other externally at A.

If we consider T the transverse, instead of the direct common tangent of O and N, then (2) becomes

$$d^2 = T_1^2 + (R + r)^2.$$

Substitute in (5) as before and


$$T_1 = 2\left((n^2 - 1)Rr\right)^{\frac{1}{2}} = \sqrt{(n^2 - 1)d_1d_n}.$$

In this case, if  $n = 2$ ,  $T_1^2 = 3d_1d_n$ .

### ~~~~~ SOLUTION OF A PROBLEM.

BY E. B. SEITZ, GREENVILLE, OHIO.

Three circles whose radii are  $a$ ,  $b$ ,  $c$ , touch each other externally. Within the space enclosed by them a circle is drawn tangent to the three circles, and within this circle three circles are drawn tangent to each other and to the three given circles. Calling the radii of these three circles  $x_1$ ,  $y_1$ ,  $z_1$ , we may determine three other circles, radii  $x_2$ ,  $y_2$ ,  $z_2$ , touching each other and the second set of circles in a similar way; and so on. Find the radii,  $x_n$ ,  $y_n$ ,  $z_n$ , of the  $n^{\text{th}}$  set of inscribed circles.



**SOLUTION.**—Let A, B, C be the centers of the given circles, GHK the circle which touches them, O its center, A', B', C' the centers of the first set of inscribed circles.

Put  $AD = a$ ,  $BD = b$ ,  $CF = c$ ,  $OG = r_1$ ,  $A'G = x_1$ ,  $B'H = y_1$ ,  $C'K = z_1$ ,  $r_0$  = the radius of the circle circumscribing the three given circles; also put  $\angle BAC = \beta$ ,

$\angle BAO = \theta$ ,  $\angle CAO = \psi$ ,  $\angle AOB = \omega$ .